

# fact sheet



Ancare Australia Pty Ltd  
 Level 6, 79 George St  
 Parramatta NSW 2150  
 Ph (02) 9893 0071  
 Fax (02) 9893 0099  
 Email:  
 genesis@ancare.com.au  
 Website:  
 www.ancare.com.au

## CYRAZIN Spray On SHEEP BLOWFLY TREATMENT

A ready-to-use spray containing 60g/L CYROMAZINE, an Insect Growth Regulator (IGR)

- Persistent activity – up to 11 weeks protection from blowfly strike
- Controls organophosphate resistant strains
- For use on long wool sheep
- Direct spraying
- Coloured with a scourable dye for identification of treated sheep

### PERSISTENT ACTIVITY

CYRAZIN prevents the development of the sheep blowfly (*Lucilia cuprina*) by interrupting the moulting of the larvae into the next developmental stage. CYRAZIN Spray On protects long wool sheep from strike for up to 11 weeks.

### CONTROLS ORGANOPHOSPHATE RESISTANT STRAINS

CYRAZIN is effective against strains of sheep blowfly that are resistant to organophosphate control products.

### FOR USE ON LONG WOOL SHEEP

CYRAZIN should only be used on sheep with at least 6 weeks of wool growth (long wool). Application rate should be calculated from the table below.

FOR THE PREVENTION OF BODY STRIKE				
Months since shearing	Total volume/sheep (mL)	Volume per treatment band (mL)	No. of bands	Number of sheep treated per 20 L
Up to 6	34	17	2	588
7	38	19	2	524
8	44	22	2	452
9	50	25	2	400
10	56	28	2	356
FOR THE PREVENTION OF BODY STRIKE AND CRUTCH STRIKE				
Months since shearing	Total volume/sheep (mL)	Volume per treatment band (mL)	No. of bands	Number of sheep treated per 20 L
Up to 6	51	17	3	392
7	57	19	3	348
8	66	22	3	300
9	75	25	3	264
10	84	28	3	236
FOR THE PREVENTION OF POLL STRIKE				
Rams and sheep susceptible to poll strike should be treated. Apply at least 5mL of CYRAZIN Spray On to the areas requiring protection, around the poll and horns.				




## DIRECT SPRAYING

**CYRAZIN Spray On** should not be diluted prior to application. It should be applied using an appropriate applicator gun, with the gun held approximately 25cm above the fleece so that a 15cm wide band is applied. Application sites to prevent body and crutch strike are illustrated in the diagrams below.



For the prevention of body strike



For the prevention of crutch strike

## COLOURED WITH A SCOURABLE DYE FOR IDENTIFICATION OF TREATED SHEEP

**CYRAZIN Spray On** contains a scourable dye so that it is easy to identify which sheep have been treated and confirm that the product has been applied in the correct areas.

## WITHHOLDING PERIODS

**MEAT:** 7 days

**EXPORT SLAUGHTER INTERVAL (ESI)\*:** 28 days

**MILK:** DO NOT USE on ewes which are producing or may in the future produce milk or milk products for human consumption.

**WOOL:** 2 months

## PACK SIZES AVAILABLE

20L Drum



\*Check APVMA website for current ESI - <http://www.apvma.gov.au/residues/ESI.shtml>