

fact sheet



Ancare Australia Pty Ltd
 Level 6, 79 George Street
 Parramatta NSW 2150
 Ph (02) 9893 0071
 Fax (02) 9893 0099
 Website:
www.ancare.com.au

EXIT POUR-ON LOUSICIDE FOR SHEEP

A ready-to-use off-shears pour-on for sheep containing 25 g/L TRIFLUMURON, an Insect Growth Regulator (IGR)

- Control of Body Lice
- Persistent activity
- Flexible application times
- Convenient backline application
- Rainfast
- Wide Safety Margin
- Short 4 week quarantine period
- Short 14 day meat withholding period

CONTROL OF BODY LICE

EXIT Pour-on controls body lice (*Bovicola ovis*) on shorn sheep up to 7 days off-shears and lambs at foot (up to 3 months of age).

EXIT Pour-on is effective against all strains of lice, including those resistant to synthetic pyrethroids.

PERSISTENT ACTIVITY

EXIT Pour-on:

- protects against the establishment of lice populations if treated sheep are exposed to re-infestation within 12 weeks of treatment;
- prevents the development of immature lice present in the fleece at application and those that hatch from eggs in the following 20 weeks; and
- produces a rapid decline in lice numbers with all adult lice present at application dying by 14 weeks.

FLEXIBLE APPLICATION TIMES

This product can be used on all sheep up to 7 days off-shears and on unshorn lambs up to 3 months of age.

When shearing ewes with lambs at foot (up to 3 months of age), shear and treat the ewes, and treat the unshorn lambs at the same time.

When lambing off-the-board, treat the ewes at shearing, and treat the unshorn lambs at any time up to 3 months of age.

EXIT Pour-on may be used in conjunction with worm control drenches and at the same time as vaccination.



CONVENIENT BACKLINE APPLICATION

Sheep are treated with a single backline application.

EXIT Pour-on contains a blue scourable dye to identify treated sheep.

RAIN FAST

Sheep should be treated dry, but moderate rainfall even within 4 days of treatment will not reduce the product's efficacy.

APPLICATION

EXIT Pour-on is applied in a single strip down the middle of the back between the poll and the butt of the tail using an applicator gun and nozzle. Dose volume should be calculated from the table below.

Bodyweight	Dose Volume	20L pack treats (head)
Up to 10kg	10 mL	2000
11 – 20 kg	12 mL	1666
21 – 30kg	15 mL	1333
31 – 55kg	20 mL	1000
56 – 75kg	25 mL	800
76 – 85kg	30 mL	666
86 – 95kg	35 mL	571

WIDE SAFETY MARGIN

EXIT Pour-on may be used in sheep of all ages. Ewes may be treated at any stage of pregnancy or during lactation.

SHORT 4 WEEK QUARANTINE PERIOD

After just 4 weeks, adult lice on treated sheep cannot infect clean sheep and treated sheep can be mixed with louse-free sheep.

WITHHOLDING PERIODS

MEAT: 14 days The short withholding period of **EXIT Pour-on** allows flexible management of lambs, including time to slaughter. This provides good capacity to respond to market fluctuations.

EXPORT SLAUGHTER INTERVAL (ESI): 66 days

MILK: DO NOT USE on ewes which are producing or may in the future produce milk or milk products for human consumption.

PACK SIZES AVAILABLE

20L Drum

